

FILE COPY FOREST INSECT LABORATORY,
STANFORD UNIVERSITY CALIFORNIA.

4541

#602

SUSCEPTIBILITY OF WESTERN YELLOW PINES

TO BARKBEETLE ATTACK

by

F. P. KEEN
Associate Entomologist
Bureau of Entomology

Forest Insect Laboratory
Stanford University, California
February, 1928

SUSCEPTIBILITY OF WESTERN YELLOW PINES TO BARKBEETLE ATTACK

by

E. P. KENN

Associate Entomologist

Office of Forest Insect Investigations, Bureau of Entomology,
United States Department of Agriculture

Does the western pine beetle attack and kill trees indiscriminately, or does it show a decided preference for certain types of yellow pines? The answer to this question has a decidedly practical application, viz.: if the type of trees susceptible to beetle attack is known, they can be removed during selective logging and the subsequent loss from this source reduced. Further, knowing the type of trees selected by beetles for their attack, any given tract of timber can be judged as to the risk of heavy beetle losses occurring in it, simply by estimating the proportion of beetle-susceptible trees in the stand. This question has puzzled foresters and forest entomologists for some time, and only recently has considerable light been thrown upon the subject.

Person, in 1926, made an intensive study of an area of about four sections near Northfork, California, and found that "the loss from D.b. per unit area varied inversely with the quality of the site, the heaviest losses being found on the poorest sites; that there appeared to be no well-defined relation between shape of crown and selection by the beetles except that sharp-pointed trees were avoided and flat-topped trees preferred to some extent; that a decided preference was shown for trees between 20 and 30" d.b.h., and that the beetles selected the more slowly-growing trees in 75 per cent of the cases studied."

The surveys on the Southern Oregon-Northern California Project, which started in 1921, have accumulated considerable data on the size, growth rate and crown classes of trees killed by barkbeetles--mainly the western pine beetle (Dendroctonus brevicornis Lec.). Each year some 50 permanent sample plots, mostly of 640 acres each and scattered throughout about 2000 square miles of the yellow pine stands of the east side type, have been intensively cruised. A three-man crew, consisting of two spotters and a compassman, run two strips through each forty. The spotters, working in a five-chain strip on each side of the compassman, cover 100 per cent of the plot and mark up all the current year's beetle loss. During the survey of 1927, 47 plots totaling 19,680 acres were covered in this manner, and in addition to the usual data all the trees killed by beetles during 1926 were tallied as to their tree class. In addition, on each plot the compassman tallied all the green trees falling within a 13.2-foot strip (2 per cent estimate) along his line, and tabulated them as to tree class and diameter. With the completion of this survey we have a sample consisting of 10,053 green trees and 13,160 beetle-killed trees, classified as to tree class and diameter and representing a wide range of conditions as to site, exposure, soil and elevation. With such a basis we have a very good index as to the class of trees selected by the western pine beetle in the timber type of this region.

TREE CLASSIFICATION

Early in 1927, Duncan Durning of the California Forest Experiment Station proposed a tree classification for western yellow pine based upon age, degree of dominance, crown development and thrift. Taking the sum of these characteristics, seven classes were defined as follows:

CLASS 1 Age class, young or thrifty mature; position, isolated or dominant (rarely co-dominant); crown length, 65 per cent or more of total height; crown width, average or wider; form of top, pointed; vigor, good (pl. I, fig. 1).

CLASS 2 Age class, young or thrifty mature; position, usually co-dominant (rarely isolated or dominant); crown length, less than 65 per cent of the total height; crown width, average or narrower; form of top, pointed; vigor, good or moderate (pl. I, fig. 2).

CLASS 3 Age class, mature; position, isolated or dominant (rarely co-dominant); crown length, 65 per cent or more of total height; crown width, average or wider; form of top, round; vigor, moderate (pl. I, fig. 3).

CLASS 4 Age class, mature; position, usually co-dominant (rarely isolated or dominant); crown length, less than 65 per cent of total height; crown width, average or narrower; form of top, round; vigor, moderate or poor (pl. I, fig. 4).

CLASS 5 Age class, overmature; position, isolated or dominant (rarely co-dominant); crown, of any size; form of top, flat; vigor, poor (pl. I, fig. 5).

CLASS 6 Age class, young or thrifty mature; position, intermediate or suppressed; crown, of any size, usually small; form of top, round or pointed; vigor, moderate or poor (pl. I, fig. 6).

CLASS 7 Age class, mature or overmature; position, intermediate or suppressed; crown, of any size, usually small; form of top, flat; vigor, poor (pl. I, fig. 7).

TREE CLASSES SELECTED

Classifying all the trees killed by beetles in 1926 and 2 per cent of the green trees on all the sample plots, we find that the average distribution of the trees in these classes within the stand is as shown in Plate II. Throughout southern Oregon and northern California, in the east-side yellow-pine type, overmature pines of Class 5 form over 32 per cent of the stand; vigorous trees of Class 1 are the next in order of abundance and represent 22 per cent of the stand; Class 3 is next in order; and the smallest number of trees, representing 4 per cent of the stand, fall in Class 7.

Forty-five per cent of the beetle losses by trees occur in Class 5, twenty per cent in Class 4 and the smallest per cent in Classes 1 and 6. Naturally, since Class 5 has the trees of largest diameter, the big volume of loss occurs in trees of this class.

Taking into account the number of trees of each class in the stand and the frequency with which such trees are attacked by beetles, we have the graph shown in Plate III.

In order to determine whether certain classes of trees are more susceptible to insect attack than other classes, it is necessary to compare the frequency with which the different classes are attacked. If no selection was made by the beetles we should expect to find the distribution of beetle losses in proportion to the occurrence of the different classes within the stand. On Plate III is graphed the ratio of the beetle losses in different tree classes to the occurrence of these classes in the stand. A ratio of 1 to 1 indicates an even distribution and no selection. A ratio of less than 1 indicates avoidance, and of more than 1 a preference. This graph therefore brings out a very decided preference on the part of the beetles for trees of Classes 7, 4 and 5 in the order named, and an avoidance of trees in Classes 1, 3, 2 and 6, also in the order named. Class 7 trees are killed 2.25 times as frequently as they should be if an even distribution of losses occurred, and Class 1 trees only one-quarter as often.

It was quite a surprise to find such a preference shown for Class 7 trees, since beetle-killed trees of this class are not frequently encountered. But this is simply because the green trees of this class form only 4 per cent of the stand, and hence are inconspicuous in numbers.

The fact that Class 4 trees are more frequently killed than Class 5 is another interesting point that is contrary to the impression secured from general observation. However, it is entirely logical that it should be so, since the mature co-dominant trees of this class are the ones suffering from the greatest competition, while trees of Class 5 are usually isolated dominants which have secured a good foothold and have plenty of ground space to draw upon for their moisture supply.

The trees of these three susceptible classes, according to studies by Dunning, all show annual growth rates of less than 1 per cent of the basal area, even on the better sites, which ties in well with other studies as to the preference of the beetles for slow-growing trees. Some interesting relationships between the volumes of beetle-killed trees and trees in the stand are brought out in Plate IV.

On comparing the distribution of the board-foot volumes in the seven tree classes, it is found that on the check plots of Area 1 over 80 per cent of the volume occurs in the overmature dominants of Class 5. On Area 2, 65 per cent of the volume is found in this class (pl. IV, Figs. 1 and 4), and on both areas over 80 per cent of the volume of beetle loss occurs in this class. The other classes represent a comparatively small proportion of the volume.

However, on comparing the frequency of the beetle losses by volume in the different tree classes with the occurrence of these volumes in the classes for the stand as a whole, we find the same selection of tree classes as is shown in Plate III, when this comparison was made on the tree basis. Again we find that Class 7 is the greatest risk, Class 4 the next in importance, while only a slight preference is shown for Class 5 trees.

On comparing the average volume of live and killed trees by classes, another suggestive point is brought out (pl. IV, fig. 3). Classes 2, 4, 6 and 7, which include all the co-dominant, intermediate and suppressed trees, show almost the same average volume for beetle-killed as for green trees. This indicates that within these classes no particular selection of large or small individuals is shown by the beetles. However, for the dominant trees of Classes 1, 3 and 5, the average volume of the killed trees is uniformly lower than that of the stand as a whole; which means that when the beetles do kill trees of these dominant classes they select the undersized individuals, which are probably less resistant than the average of their class. This point is most significant.

In Plate V, this same ratio of occurrence of tree classes in insect losses to the occurrence in the stand is shown for the four areas of the Southern Oregon-Northern California region. In every case practically the same ratios hold as for the combined results, which shows that the basis is an entirely adequate one for definite conclusions and that the same selective habits of the beetles hold throughout this region.

However, we note some slight variations on these different areas. For instance, on Area 2, Class 4 trees are not so prominently susceptible as in the general case; on Area 3, Class 2 trees show some slight degree of susceptibility, and on the Modoc area Class 7 trees are outranked by Class 4. To determine if these variations are due to different site conditions the data have been rearranged according to site.

SELECTION OF TREE CLASSES ON VARIOUS SITES

Plate VI shows the distribution of green trees and beetle-killed trees on Sites 2, 3, 4 and 5. No plot running to Site 1 was available on the area. This chart shows about the same distribution of trees by tree classes on all sites, without any striking differences. But when we compare the ratio of beetle-killed trees to the total trees in the stand on the different sites, some striking differences are brought out.

Thus in Plate VII on Site 2 we find a very pronounced selection of 4's and 7's, the co-dominant, intermediate and suppressed trees of the mature and overmature age classes. Overmature trees of Class 5 on this site are able to maintain themselves, and hence are only slightly susceptible.

On Site 3 we still have a strong preference shown for 4's and 7's. On Site 4, which forms the bulk of the pine stand in this region, we have the same preference for 4's, 5's and 7's as was brought out in Plate III for the stand as a whole. On Site 5 we find that Class 5 has a hard struggle, and the preference of the beetles is about equally divided between 4's, 5's and 7's.

Using these same figures and plotting the trees of each class according to site and per cent of stand killed, certain interesting sidelights are brought out (Plate VIII).

Trees of Class 1 form from 18.1 per cent to 24.6 per cent of the stand on the various sites. They form a higher per cent of the stand on poor sites than on good ones, showing that on good sites most of the other and older tree classes survive to form a large part of the stand, while on poor sites the older trees are more frequently killed, with a higher proportion of Class 1 trees surviving. Only from 1.7 to 5.4 per cent of the beetle loss is found in trees of this class, with a higher percentage of such trees killed on the poor sites than on the good ones.

Class 2 trees form from 9.1 to 12.8 per cent of the stand, and in beetle losses they represent from 6.7 to 9.7 per cent of the total loss. There does not appear to be any marked difference in the occurrence of such trees or the losses by sites.

Trees of Class 3 show the same tendency as trees of Class 1. As the site gets poorer they form a larger percentage of the stand, due to the greater elimination of less dominant trees. The beetle losses also increase in this class as the site gets poorer, but still do not occur as frequently as do the trees in the stand.

For trees of Class 4, we see that on every site they appear more frequently in the beetle losses than they do in the stand as a whole. However, as the site gets poorer the proportion of Class 4 trees consistently decreases. This is because on the better sites there is a greater difference in the growth rate, and thus of susceptibility, than on the poorer sites. On the latter all classes of trees tend to become susceptible, and thus a smaller percentage of the losses fall in any particular tree class.

For trees of Class 5, the beetle losses in every case appear more frequently than these trees occur in the stand, and the ratio of loss increases steadily as the site becomes poorer. The percentage of green trees in this class varies from 33.1 per cent on Site 2 to 29.0 per cent on Site 5. This indicates that the overmature trees of this class maintain themselves longer on a good site than they do on a poor one, and hence form a higher percentage of the total stand. As would be expected, the beetle losses are increasingly greater on the poorer sites.

Trees of Class 6 represent only a very small percentage of the stand, varying from 7.2 per cent on Site 2 to 5.5 per cent on Site 5. The beetles do not show any particular preference for these trees and kill a varying proportion on the different sites.

Trees of Class 7 also form only a small percentage of the stand, running around 4 to 5 per cent for all sites. The beetle losses in every case occur with greater frequency in this class than they would if an even distribution of loss occurred. The competition and hence beetle loss appear to be greater for trees on the good sites than on the poor ones.

SELECTION OF TREES BY DIAMETER GROUPS

The samples of green trees and all the beetle-killed trees of the 1926 loss were also compared as to diameters.

Plate IX, fig. 1, shows by diameter classes the normal distribution of trees in the stand and those beetle-killed. This curve is similar in every way to the one found in the study by Person and Sanford. The big bulk of the infestation occurs in the diameter groups between 16 and 28 inches. In Fig. 2 is determined the ratio of occurrence of beetle-killed trees to that of trees in the stand for each diameter class. This graph shows a definite selection by the beetles of trees between 16 and 28 inches d.b.h., while in the diameter classes above 28 inches they kill about as frequently or less frequently than these trees occur in the stand, and hence cannot be considered as showing any preference.

Why should the beetles select trees between these diameter limits? In general it would seem reasonable that trees falling between these limits include most of the trees suffering from severe competition. The bulk of trees below 16 inches are young, thrifty and fast-growing, while the majority over 28 inches have reached a dominant position in the stand, crowded out their nearest neighbors, and are now in a better position to survive than trees of smaller diameters.

Working on this theory the tree classes of each diameter group were plotted to show the percentage of green and killed trees in each class.

Referring to Plate X, Fig. 1, we see that of trees 16 inches d.b.h., 56 per cent are young or young thrifty trees of Class 1 and 26 per cent are young co-dominants of Class 2. These two classes together form 82 per cent of the stand. All other classes are inferior trees for this diameter, and hence are the ones suffering from the greatest beetle loss.

In trees of 18 inches d.b.h., Class 1 forms 36 per cent of the stand, and 21 per cent now come in Class 3, which is beginning to be an important tree class for this diameter group. Here the bulk of the beetle loss occurs in trees that are struggling to survive--co-dominant matures, overmatures and mature suppressed. By the time the trees have reached 20 inches d.b.h., Class 3 forms the greater part of the stand, Class 4 is also prominent, and Class 1 has subsided to a little over 21; and it is in the three classes 3, 4 and 5 that the big struggle with the insects is occurring with a particularly high mortality in trees that have become overmature at this size.

In diameter groups 22 inches and 24 inches the same factors are evident, with Class 5 becoming increasingly important. With diameter 26 inches Class 5 becomes the most conspicuous class, and 67 per cent of the beetle loss falls here. By the time diameter 30 inches is reached, 67 per cent of the green trees and 86 per cent of the beetle losses occur in the overmature trees of Class 5, and for trees larger than 30 inches the beetle loss is confined almost entirely to this, since it is about the only class represented in the stand.

The evidence therefore supports the theory that the reason trees between 16 and 28 inches are selected is because the greatest struggle of the mature co-dominants is occurring in these diameter classes. After the trees reach 28 inches the bulk of trees in the stand is made up of overmature dominants that are fairly well fitted to survive.

So far all this study pertains to conditions as they existed in a year of epidemic infestation like 1926. Whether under endemic infestation conditions would be markedly different cannot well be determined until we have an endemic situation on this area. However, in 1923 the infestation was as nearly endemic as it is ever apt to be in this region. A study of the diameter groups selected by the beetles in 1923 as compared to those selected in 1926 is given in Plate XI. This shows that during the endemic year of 1923 the selection of trees between 16 and 32 inches was even more pronounced than in 1926, during the epidemic year. In both years the percentage of trees selected over 32 inches d. b.h. is about the same as the percentage of such trees in the stand, indicating that no preference was shown for these large trees, but that they were killed in about the same ratio as their occurrence in the stand.

APPLICATION OF RESULTS

Since it is evident that certain trees are more susceptible than others to barkbeetle attack, the question arises as to what will happen if the susceptible trees are removed from the forest through selective thinning. There are two alternatives: either the beetles will ignore their preferences and select supposedly less susceptible trees or they will die out because of the reduction in their normal food supply.

If the beetles disregard their preferences when a shortage of susceptible trees exists, we should expect to find them attacking all classes of trees on the better sites. Instead, however, the loss on the better sites is low and preference for the few available susceptible trees extremely well marked. If these few susceptible trees were removed we should expect to find a further reduction in the losses.

Since on the poorer sites nearly all classes of trees are susceptible, we should not expect to get such beneficial results from thinning. However, even here a thinning by reducing competition between individual trees and groups of trees should increase the vigor of the remaining stand and help to reduce the beetle losses.

The selective logging of a beetle-infested area, with removal of the more susceptible trees, should result in marked improvement in the vigor and resistance of the stand and reduce beetle losses to a minimum. However, in most of the virgin western yellow pine stands of southern Oregon and northern California, from 65 to 80 per cent of the volume of the stand is in overmature trees. The removal of susceptible trees on such areas would mean about as heavy cutting as is now practised on commercial logging operations in this region.

SUMMARY

A study of a large series of green and beetle-killed western yellow pines in southern Oregon and northern California shows that the western pine beetle in this region exhibits a decided preference for overmature, suppressed trees of Class 7, mature co-dominant trees of Class 4 and overmature dominant trees of Class 5.

The classes of trees selected are all making an annual growth of less than one per cent of the basal area.

On the areas studied, from 65 to 80 per cent of the stand and 80 per cent of the loss by volume consists of the overmature dominant trees of Class 5.

Within the co-dominant, intermediate and suppressed classes, the beetles show no selection as to size of the individuals. In the dominant classes, the average volume of trees killed by beetles is much lower than the average volume of the class as a whole.

Poor sites show a greater loss than good sites. On the poorest sites the selective tendency becomes less marked, the trees of all classes being reduced to such a low stage of vigor as to become susceptible to attack. On good sites there is a very marked selection of the suppressed and co-dominant trees of Classes 4 and 7.

As the site becomes poorer the proportion of loss increases in the dominant classes 1, 3 and 5, and decreases in the co-dominant, intermediate and suppressed classes 2, 4, 6 and 7.

The beetles show a preference for trees of the diameter classes between 16 and 32 inches d.b.h. This is principally due to the high proportion of trees struggling for dominance within this diameter range.

During endemic conditions the selection of trees within these diameter limits (16 to 32 inches) is even more marked than during epidemic conditions.

The results of this study emphasize the importance of marking for removal trees of classes 4, 5 and 7. It also indicates that perhaps more smaller-diameter trees in Classes 4, 5 and 7 should be removed than has heretofore been the practice. Class 3 trees are the safest risk where large trees are needed for seed or where a reserve volume of valuable material is desired. Class 2 trees are a relatively poor risk, and should be removed in preference to Class 1 trees where groups are to be thinned.

The selective thinning of stands, with the removal of beetle-susceptible trees, offers hope of greatly reducing the loss from insects, especially on the better sites.

TABLE 1.-Data For Plates 2 and 3

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Total |
|-------------------------------|---------|--------|--------|--------|---------|--------|--------|---------|
| Green Trees on Strips | 2,227 | 1,003 | 1,514 | 1,154 | 3,147 | 605 | 403 | 10,053 |
| Total Green on Plots | 111,350 | 50,150 | 75,700 | 57,700 | 157,350 | 30,250 | 20,150 | 502,650 |
| Beetle-Killed | 657 | 913 | 1,256 | 2,567 | 5,930 | 652 | 1,185 | 13,160 |
| Total | 112,007 | 51,063 | 76,956 | 60,267 | 163,280 | 30,902 | 21,335 | 515,810 |
| Per cent Occurrence | 21.7 | 9.9 | 14.9 | 11.7 | 31.7 | 6.0 | 4.1 | 100% |
| Per cent Occurrence of Killed | 5.0 | 6.9 | 9.5 | 19.5 | 45.0 | 4.9 | 9.0 | 100% |
| Per cent of Stand Killed | .59 | 1.73 | 1.63 | 4.26 | 3.63 | 2.11 | 5.55 | 2.55 |
| Ratio of Occurrence | .23 | .70 | .64 | 1.67 | 1.42 | .82 | 2.20 | 1 |

TABLE 2.—Data for Plate 5

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Total |
|-------------------------------|--------|--------|--------|--------|--------|-------|-------|---------|
| Green Trees in Strips | 583 | 343 | 293 | 363 | 1,141 | 127 | 132 | 2,982 |
| Green Trees on Plots | 29,150 | 17,150 | 14,650 | 13,150 | 57,050 | 6,350 | 6,600 | 149,100 |
| Beetle-Killed (1926) | 79 | 100 | 125 | 438 | 871 | 19 | 192 | 1,924 |
| Total | 29,229 | 17,250 | 14,775 | 13,588 | 57,921 | 6,369 | 6,792 | 150,924 |
| Per cent Occurrence in Stand | 19.4 | 11.4 | 9.8 | 12.3 | 38.4 | 4.2 | 4.5 | 100% |
| Per cent Occurred Killed 1926 | 4.3 | 5.5 | 6.9 | 24.0 | 47.8 | 1.0 | 10.5 | 100% |
| Ratio | .22 | .43 | .70 | 1.95 | 1.24 | .24 | 2.34 | 1 |

AREA 2

| | | | | | | | | |
|-----------------|--------|--------|--------|--------|--------|-------|-------|---------|
| Green | 447 | 196 | 493 | 332 | 585 | 162 | 105 | 2,320 |
| Green on Plots | 22,350 | 9,800 | 24,650 | 16,600 | 29,250 | 8,100 | 5,280 | 116,000 |
| Beetle-Killed | 309 | 231 | 598 | 918 | 2,371 | 240 | 500 | 5,167 |
| Total | 22,659 | 10,031 | 25,248 | 17,518 | 31,621 | 8,340 | 5,780 | 121,167 |
| Per cent Stand | 18.7 | 8.3 | 20.8 | 14.5 | 26.1 | 6.9 | 4.7 | 100% |
| Per cent Killed | 6.0 | 4.5 | 11.5 | 17.8 | 45.8 | 4.6 | 9.7 | 100% |
| Ratio | .32 | .54 | .56 | 1.23 | 1.75 | .67 | 2.06 | 1 |

AREA 3

| | | | | | | | | |
|-----------------|--------|--------|--------|--------|--------|--------|-------|---------|
| Green Strip | 734 | 303 | 564 | 333 | 1,016 | 244 | 110 | 3,304 |
| Plot | 36,700 | 15,150 | 29,200 | 16,650 | 50,800 | 12,200 | 5,500 | 165,200 |
| Killed | 241 | 515 | 430 | 904 | 2,167 | 349 | 400 | 5,086 |
| Total | 36,941 | 15,665 | 29,630 | 17,554 | 52,967 | 12,549 | 5,900 | 170,286 |
| Per cent Stand | 21.6 | 9.2 | 16.8 | 10.4 | 31.1 | 7.4 | 3.5 | 100% |
| Per cent Killed | 4.7 | 10.1 | 8.4 | 19.3 | 42.7 | 6.9 | 7.9 | 100% |
| Ratio | .22 | 1.10 | .50 | 1.86 | 1.37 | .93 | 2.26 | 1 |

HODOC

| | | | | | | | | |
|-----------------|--------|-------|-------|-------|--------|-------|-------|--------|
| Green Strip | 453 | 161 | 164 | 126 | 405 | 72 | 56 | 1,447 |
| Green on Plots | 23,150 | 8,050 | 8,200 | 6,800 | 20,250 | 3,600 | 2,800 | 72,350 |
| Killed | 28 | 67 | 103 | 227 | 521 | 44 | 93 | 1,083 |
| Total | 23,178 | 8,117 | 8,303 | 6,927 | 20,771 | 3,644 | 2,893 | 73,433 |
| Per cent Stand | 31.5 | 11.1 | 11.3 | 8.9 | 28.3 | 5.0 | 3.9 | 100% |
| Per cent killed | 2.6 | 6.2 | 9.5 | 21.0 | 49.1 | 4.0 | 8.6 | 100% |
| Ratio | .08 | .56 | .84 | 2.36 | 1.70 | .80 | 2.21 | 1 |

TABLE 3.—Data for Plates 6, 7 and 8

SITE II

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Total |
|-----------------------|-------|-------|-------|-------|--------|-------|-------|--------|
| Green Trees on Strips | 138 | 88 | 58 | 63 | 260 | 49 | 27 | 683 |
| Total Green on Plots | 6,900 | 4,400 | 2,900 | 3,150 | 13,000 | 2,450 | 1,350 | 34,150 |
| Beetle-Killed | 3 | 17 | 4 | 48 | 72 | 12 | 20 | 176 |
| Total | 6,903 | 4,417 | 2,904 | 3,198 | 13,072 | 2,462 | 1,370 | 34,326 |
| Per cent Stand | 20.1 | 12.8 | 8.5 | 9.3 | 38.1 | 7.2 | 4.0 | 100% |
| Per cent Killed | 1.7 | 9.7 | 2.3 | 27.2 | 43.9 | 6.8 | 11.4 | 100% |
| Ratio | .08 | .76 | .27 | 2.32 | 1.07 | .95 | 2.85 | |

SITE III

| | | | | | | | | |
|-----------------------|--------|--------|-------|--------|--------|-------|-------|--------|
| Green Trees on Strips | 311 | 203 | 161 | 226 | 616 | 103 | 85 | 1,703 |
| Total Green on Plots | 15,550 | 10,150 | 8,050 | 11,300 | 30,800 | 5,150 | 4,150 | 85,150 |
| Beetle-Killed | 14 | 50 | 49 | 186 | 305 | 17 | 124 | 745 |
| Total | 15,564 | 10,200 | 8,099 | 11,486 | 31,105 | 5,167 | 4,274 | 85,895 |
| Per cent Stand | 18.1 | 11.9 | 9.4 | 13.4 | 36.2 | 6.0 | 5.0 | 100% |
| Per cent Killed | 1.9 | 6.7 | 6.6 | 24.9 | 41.0 | 2.3 | 16.6 | 100% |
| Ratio | .10 | .56 | .70 | 1.86 | 1.13 | .38 | 3.32 | |

SITE IV

| | | | | | | | | |
|-----------------------|--------|--------|--------|--------|---------|--------|--------|---------|
| Green Trees on Strips | 1,669 | 668 | 1,218 | 820 | 2,151 | 438 | 276 | 7,240 |
| Total Green on Plots | 83,450 | 53,400 | 60,900 | 41,000 | 107,550 | 21,900 | 13,800 | 362,000 |
| Beetle-Killed | 616 | 778 | 1,095 | 2,174 | 5,100 | 593 | 978 | 11,334 |
| Total | 84,066 | 54,178 | 61,995 | 43,174 | 112,650 | 22,493 | 14,778 | 373,334 |
| Per cent Stand | 22.5 | 9.1 | 16.6 | 11.6 | 30.2 | 6.0 | 4.0 | 100% |
| Per cent Killed | 5.4 | 6.9 | 9.7 | 19.2 | 45.0 | 5.2 | 8.6 | 100% |
| Ratio | .24 | .76 | .58 | 1.66 | 1.49 | .87 | 2.15 | |

SITE V

| | | | | | | | | |
|-----------------------|-------|-------|-------|-------|-------|-----|------|--------|
| Green Trees on Strips | 109 | 44 | 77 | 45 | 120 | 15 | 17 | 427 |
| Total Green on Plots | 5,450 | 2,200 | 3,850 | 2,250 | 6,000 | 750 | 850 | 21,350 |
| Beetle-Killed | 24 | 68 | 108 | 159 | 453 | 30 | 63 | 905 |
| Total | 5,474 | 2,268 | 3,958 | 2,409 | 6,453 | 780 | 913 | 22,255 |
| Per cent Stand | 24.6 | 10.2 | 17.8 | 10.8 | 29.0 | 3.5 | 4.1 | 100% |
| Per cent Killed | 2.6 | 7.5 | 12.0 | 17.6 | 50.0 | 3.3 | 7.0 | 100% |
| Ratio | .10 | .74 | .67 | 1.63 | 1.72 | .94 | 1.71 | |

TABLE 4.--Data for Plate IX

Distribution of Trees by Diameter Classes

| | 10 | 12 | 14 | 16 | 18 | 20 | 22 | 24 | 26 | 28 | 30 | 32 | 34 | 36 | 38 | 40 | 42 | 44 | 46 | 48 | 50 | 52 | 54 | 56 | 58 | 60 | 62 | 64 | 66 | | |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|--------|-------|------|------|------|------|------|------|------|---------------|---------|
| Green | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
| Strip | 893: | 926: | 840: | 666: | 778: | 737: | 744: | 737: | 696: | 702: | 625: | 481: | 366: | 292: | 229: | 144: | 83: | 59: | 41: | 47: | 32: | 15: | 13: | 14: | 9: | 4: | 2: | 1: | 1: | : | |
| Green | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
| Plots | 44,650: | 46,300: | 42,000: | 33,300: | 38,900: | 36,850: | 37,200: | 36,850: | 34,800: | 35,100: | 31,250: | 24,050: | 19,300: | 14,600: | 11,450: | 7,200: | 4,150: | 2,950: | 2,050: | 2,350: | 1,600: | 750: | 650: | 700: | 450: | 200: | 100: | 50: | 50: | Total Green | 509,850 |
| Killed | 688: | 1,044: | 1,085: | 1,047: | 1,203: | 1,239: | 1,196: | 1,512: | 1,251: | 1,015: | 698: | 641: | 523: | 263: | 238: | 134: | 110: | 88: | 57: | 41: | 30: | 21: | 15: | 6: | 2: | 4: | 2: | 0: | 0: | Total Killed | 14,153 |
| Total | 45,338: | 47,344: | 43,085: | 34,347: | 40,103: | 38,089: | 38,396: | 38,362: | 36,051: | 36,115: | 31,948: | 24,691: | 19,823: | 14,863: | 11,688: | 7,334: | 4,260: | 3,038: | 2,107: | 2,391: | 1,630: | 771: | 665: | 706: | 452: | 204: | 102: | 50: | 50: | Total on Plot | 524,003 |
| Per ct. | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
| Stand | 8.65: | 9.03: | 8.21: | 6.55: | 7.64: | 7.26: | 7.33: | 7.32: | 6.88: | 6.89: | 6.10: | 4.72: | 3.79: | 2.84: | 2.24: | 1.40: | .91: | .58: | .41: | .46: | .31: | .15: | .13: | .13: | .09: | .04: | .02: | .01: | .01: | : | |
| Per ct. | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
| Killed | 4.86: | 7.38: | 7.66: | 7.39: | 8.31: | 8.75: | 8.45: | 10.71: | 8.86: | 7.18: | 4.94: | 4.54: | 3.69: | 1.86: | 1.68: | .96: | .78: | .62: | .41: | .30: | .22: | .15: | .10: | : | : | : | : | : | : | : | |
| Ratio | .56: | .82: | .93: | 1.13: | 1.11: | 1.21: | 1.15: | 1.46: | 1.29: | 1.04: | .81: | .96: | .98: | .66: | .75: | .63: | .96: | 1.07: | 1.00: | .65: | .71: | 1.00: | .77: | : | : | : | : | : | : | : | |

TABLE 5.—Data for Plate X
Tree Classes

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Total |
|-----------------|---------|--------|---------|---------|---------|--------|--------|--------|
| Diam. 16" | : | : | : | : | : | : | : | : |
| Green Strips | 372: | 170: | 1: | 4: | 0: | 21: | 75: | 643 |
| Green Plots | 18,600: | 8,500: | 50: | 200: | 0: | 1050: | 3,750: | 32,150 |
| Beetle-Killed | 77: | 170: | 129: | 248: | 80: | 50: | 233: | 987 |
| Total | 18,677: | 8,670: | 179: | 448: | 80: | 1,100: | 3,983: | 33,157 |
| Per cent Stand | 56.4: | 26.2: | .5: | 1.4: | .2: | 3.3: | 12.0: | 100% |
| Per cent Killed | 7.7: | 17.2: | 13.1: | 25.2: | 8.1: | 5.1: | 23.6: | 100% |
| Diam. 18" | : | : | : | : | : | : | : | : |
| Green Strips | 284: | 108: | 162: | 135: | 4: | 4: | 59: | 756 |
| Green Plots | 14,200: | 5,400: | 8,100: | 6,750: | 200: | 200: | 2,950: | 37,800 |
| Beetle-Killed | 65: | 153: | 165: | 360: | 215: | 30: | 160: | 1,148 |
| Total | 14,265: | 5,553: | 8,265: | 7,110: | 415: | 230: | 3,110: | 33,948 |
| Per cent Stand | 36.6: | 14.3: | 21.2: | 18.2: | 1.1: | .6: | 8.0: | 100% |
| Per cent Killed | 5.7: | 13.3: | 14.4: | 31.3: | 18.7: | 2.6: | 14.0: | 100% |
| Diam. 20" | : | : | : | : | : | : | : | : |
| Green Strips | 160: | 52: | 222: | 213: | 82: | 0: | 3: | 732 |
| Green Plots | 8,000: | 2,600: | 11,100: | 10,650: | 4,100: | 0: | 150: | 36,600 |
| Beetle-Killed | 38: | 64: | 188: | 410: | 452: | 11: | 34: | 1,197 |
| Total | 8,038: | 2,664: | 11,288: | 11,060: | 4,552: | 11: | 184: | 37,797 |
| Per cent Stand | 21.3: | 7.1: | 29.8: | 29.3: | 12.0: | .0: | .5: | 100% |
| Per cent Killed | 3.2: | 5.4: | 15.7: | 34.2: | 37.8: | .9: | 2.8: | 100% |
| Diam. 22" | : | : | : | : | : | : | : | : |
| Green Strips | 87: | 20: | 228: | 229: | 155: | 0: | 0: | 719 |
| Green Plots | 4,330: | 1,000: | 11,400: | 11,450: | 7,750: | 0: | 0: | 35,950 |
| Beetle-Killed | 21: | 48: | 141: | 399: | 491: | 17: | 13: | 1,130 |
| Total | 4,371: | 1,048: | 11,541: | 11,849: | 8,241: | 17: | 13: | 37,080 |
| Per cent Stand | 11.8: | 2.8: | 31.1: | 32.1: | 22.2: | 0: | 0: | 100% |
| Per cent Killed | 1.8: | 4.2: | 12.5: | 35.4: | 43.5: | 1.5: | 1.1: | 100% |
| Diam. 24" | 1 | 2 | 3 | 4 | 5 | 6 | 7 | Total |
| Green Strips | 64: | 6: | 246: | 199: | 201: | 0: | 0: | 716 |
| Green Plots | 3,200: | 300: | 12,300: | 9,950: | 10,050: | 0: | 0: | 35,800 |
| Beetle-Killed | 15: | 17: | 144: | 449: | 745: | 57: | 7: | 1,434 |
| Total | 3,215: | 317: | 12,444: | 10,399: | 10,795: | 57: | 7: | 37,234 |
| Per cent Stand | 8.7 | .8 | 33.4 | 27.9 | 29.0 | .2: | 0: | 100% |
| Per cent Killed | 1.0 | 1.2 | 10.0 | 31.4 | 51.9 | 4.0: | .5: | 100% |
| Diam. 26" | : | : | : | : | : | : | : | : |
| Green Strips | 44: | 4: | 214: | 143: | 274: | : | : | 673 |
| Green Plots | 2,200: | 200: | 10,650: | 7,150: | 13,700: | : | : | 33,900 |
| Beetle-Killed | 4: | 7: | 99: | 245: | 791: | 30: | 4: | 1,180 |
| Total | 2,204: | 207: | 10,749: | 7,395: | 14,491: | 30: | 4: | 35,080 |
| Per cent Stand | 6.3 | .6 | 30.6 | 21.1 | 41.3 | .1: | 0: | 100% |
| Per cent Killed | .3 | .6 | 8.4 | 20.8 | 67.1 | 2.5: | .3: | 100% |
| Diam. 28" | : | : | : | : | : | : | : | : |
| Green Strips | 32: | 2: | 176: | 105: | 382: | 0: | 0: | 697 |
| Green Plots | 1,600: | 100: | 8,800: | 5,350: | 19,100: | 0: | 0: | 34,850 |
| Beetle-Killed | 6: | 3: | 63: | 159: | 729: | 0: | 0: | 960 |
| Total | 1,606: | 103: | 8,863: | 5,409: | 19,829: | 0: | 0: | 35,810 |
| Per cent Stand | 4.4 | .3 | 24.8 | 15.1 | 55.4 | 0: | 0: | 100% |
| Per cent Killed | .6 | .3 | 6.6 | 15.6 | 75.9 | 0: | 0: | 100% |
| Diam. 30" | : | : | : | : | : | : | : | : |
| Green Strips | 27: | 0: | 112: | 64: | 417: | 0: | 0: | 620 |
| Green Plots | 1,350: | 0: | 5,600: | 3,200: | 20,850: | 0: | 0: | 31,000 |
| Beetle-Killed | 2: | 1: | 30: | 53: | 553: | 0: | 0: | 639 |
| Total | 1,352: | 1: | 5,630: | 3,253: | 21,403: | 0: | 0: | 31,639 |
| Present Stand | 4.3 | 0: | 17.8 | 10.3 | 67.6 | 0: | 0: | 100% |
| Per cent Killed | .3 | .2 | 4.7 | 8.3 | 86.5 | 0: | 0: | 100% |

TABLE 6.—Data for Plate IV
Green

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | | | | | | | |
|-------------|--------|-----------|--------|---------|--------|------------|--------|-----------|--------|------------|--------|---------|-----|--------|
| Unit: | Trees: | Volume: | Trees: | Volume: | Trees: | Volume: | Trees: | Volume: | Trees: | Volume: | Trees: | Volume: | | |
| Antelope28: | 53: | 508: | 35: | 256: | 45: | 20,64: | 21: | 719: | 92: | 11,807: | 9: | 49: | 9: | 73 |
| Ferguson33: | 44: | 295: | 19: | 113: | 49: | 16,22: | 25: | 884: | 47: | 6,727: | -: | -: | 13: | 130 |
| Gerber 24: | 64: | 575: | 14: | 110: | 65: | 26,25: | 28: | 11,04: | 84: | 93,91: | 28: | 162: | 5: | 31 |
| Goodlowe 5: | 58: | 643: | 16: | 166: | 79: | 39,84: | 52: | 22,37: | 152: | 192,28: | 41: | 255: | 18: | 118 |
| Royston 10: | 44: | 767: | 25: | 269: | 51: | 30,98: | 53: | 44,40: | 21: | 35,69: | 22: | 151: | 7: | 70 |
| Sycan 2: | 32: | 230: | 18: | 139: | 44: | 23,80: | 28: | 17,30: | 63: | 105,36: | 4: | 16: | 8: | 52 |
| Willow 20: | 66: | 928: | 13: | 168: | 67: | 42,84: | 50: | 29,69: | 88: | 113,31: | 20: | 167: | 19: | 223 |
| Total | 361: | 39,96: | 138: | 12,21: | 400: | 200,57: | 257: | 140,83: | 547: | 725,89: | 124: | 800: | 79: | 697 |
| x 50 | | 1998,00: | | 610,50: | | 10028,50: | | 7041,50: | | 36294,50: | | 400,00: | | 348,50 |
| Bugs | | 23,03: | | 25,95: | | 146,99: | | 260,55: | | 2007,12: | | 17,13: | | 38,48 |
| Tot. Stand | | 2,021,03: | | 636,45: | | 10,175,49: | | 7,302,05: | | 38,301,62: | | 417,13: | | 336,98 |
| Stand % | | 3.4: | | 1.1: | | 17.2: | | 12.3: | | 64.7: | | .7: | | .6 |
| Beetles % | | .9: | | 1.0: | | 5.8: | | 10.3: | | 79.8: | | .7: | | 1.5 |

Total, 1896 trees

59,240,750 bd.ft.

TABLE 7.--Data for Plate IV

Area 2 - Bugs

| Unit | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|--------------|------------|------------|--------------|-------------|----------------|-------------|------------|
| | Trees:Vol. | Trees:Vol. | Trees:Vol. | Trees: Vol. | Trees: Vol. | Trees: Vol. | Trees:Vol. |
| Antelope 28: | 7: 51: | 12: 75: | 19: 417: | 77: 2341: | 149: 14,491: | 6: 32: | 31: 293 |
| Ferguson 35: | 25: 207: | 10: 55: | 106: 3142: | 87: 3587: | 421: 41,700: | 7: 30: | 57: 448 |
| Gerber 24: | 9: 78: | 54: 599: | 74: 1597: | 97: 2621: | 234: 25,680: | 33: 174: | 45: 427 |
| Goodlowe 5: | 14: 82: | 27: 410: | 13: 286: | 46: 1357: | 76: 8,402: | 15: 96: | 16: 145 |
| Royston 10: | 27: 640: | 21: 318: | 36: 2204: | 77: 3709: | 54: 5,466: | 38: 338: | 29: 227 |
| Sycan 2: | 45: 266: | 15: 58: | 159: 2979: | 144: 3495: | 763: 76,343: | 28: 133: | 115: 1111 |
| Willow 20: | 71: 979: | 55: 1080: | 99: 4074: | 217: 9145: | 376: 28,630: | 76: 905: | 86: 1197 |
| Total | 198: 2303: | 192: 2595: | 506: 14,699: | 745: 26055: | 2128: 200,712: | 203: 1713: | 379: 3848 |

Total - 4351 trees

2,519,250 bd.ft.

TABLE 8.--Data for Plate IV

Area 1 - Green Trees

| | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | 7 | |
|--------------------|------|----------|------|----------|------|----------|------|----------|------|-------------|-------|----------|------|--------|
| | Trs: | Vol. | Trs: | Vol. | Trs: | Vol. | Trs: | Vol. | Trs: | Vol. | Trs: | Vol. | Trs: | Vol. |
| Clover Sta. 16: | 86: | 1,455: | 62: | 724: | 28: | 1,715: | 40: | 1,735: | 173: | 31,696: | 43: | 330: | 21: | 215 |
| Clover Sta. 36: | 95: | 1,260: | 55: | 600: | 50: | 3,148: | 66: | 3,148: | 205: | 33,243: | 37: | 286: | 31: | 357 |
| Aspen Lake 34: | 47: | 659: | 34: | 492: | 39: | 2,785: | 75: | 3,745: | 142: | 24,894: | 10: | 83: | 12: | 107 |
| Eagle Ridge 16: | 71: | 811: | 65: | 542: | 36: | 1,915: | 50: | 2,101: | 146: | 20,431: | 22: | 103: | 7: | 59 |
| Jenny Creek 33: | 46: | 380: | 38: | 255: | 27: | 971: | 39: | 1,526: | 68: | 8,086: | 7: | 41: | 13: | 107 |
| Jenny Creek 34: | 37: | 312: | 12: | 70: | 38: | 2,047: | 29: | 944: | 61: | 8,714: | 8: | 40: | 15: | 126 |
| Johnson Pra. 7: | 33: | 768: | 35: | 713: | 16: | 2,070: | 26: | 2,443: | 124: | 46,045: | -: | -: | 5: | 93 |
| Klamath Cr. 9: | 16: | 120: | 6: | 32: | 14: | 384: | 10: | 452: | 10: | 903: | 2: | 14: | 2: | 9 |
| Pokegama 36: | 43: | 350: | 17: | 148: | 20: | 1,133: | 14: | 495: | 83: | 12,835: | -: | -: | 12: | 108 |
| Round Lake 5: | 69: | 766: | 14: | 129: | 19: | 974: | 11: | 583: | 90: | 13,522: | 2: | 8: | 9: | 71 |
| | | 543: | | 6,881: | 338: | 3,705: | 235: | 17,142: | 360: | 17,173: | 1107: | 200,369: | 131: | 905: |
| | | | | | | | | | | | | | | 127: |
| Av. Volume | | 13: | | 11: | | 60: | | 48: | | 180: | | 7: | | 10 |
| Tot. Vol. on Plot: | | 344,050: | | 185,250: | | 857,100: | | 958,650: | | 10,018,450: | | 45,250: | | 63,100 |
| Bugs | | 365: | | 1,362: | | 4,038: | | 23,391: | | 135,539: | | 135: | | 2,206 |
| | | 344,415: | | 185,612: | | 861,158: | | 982,041: | | 10,153,989: | | 45,385: | | 63,306 |
| | | 2.7+: | | 1.5: | | 6.9: | | 7.0+: | | 81.0: | | .4: | | .5 |

Total - 2891 trees 12,538,906 bd.ft.

TABLE 9.—Data for Plate IV

Area 1 - Bug Trees

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|-------------|----------|--------------|--------------|--------------|---------------|----------|------------|
| | Trs:Vol: | Trs:Vol: | Trs:Vol: | Trs:Vol: | Trs:Vol: | Trs:Vol: | Trs:Vol: |
| Aspen | 34: 1: | 8: 25: 405: | 7: 306: 17: | 892: 47: | 8,335: | -: -: | 9: 115 |
| Clover | 16: 2: | 14: 9: 74: | 3: 50: 51: | 3,051: 96: | 16,459: | 5: 27: | 16: 120 |
| Clover Ste. | 36: 3: | 21: 8: 75: | 3: 95: 21: | 1,369: 58: | 9,477: | 2: 24: | 15: 160 |
| Eagle Ridge | 16: -: | -: 14: 311: | 5: 307: 53: | 1,839: 103: | 17,832: | 4: 23: | 22: 198 |
| Jenny Creek | 33: 7: | 56: 2: 24: | 27: 852: 63: | 2,183: 52: | 6,729: | 2: 16: | 10: 86 |
| Jenny Creek | 34: 8: | 66: 3: 33: | 20: 800: 94: | 3,749: 155: | 12,554: | 3: 18: | 65: 630 |
| Johnson | 7: -: | -: 1: 40: | 1: 9: 10: | 741: 44: | 8,538: | 1: 14: | 9: 189 |
| Klamath C. | 9: 10: | 137: 10: 75: | 20: 473: 36: | 1,131: 60: | 4,857: | -: -: | 11: 89 |
| Pokegama | 36: 4: | 35: 2: 13: | 17: 682: 83: | 2,475: 189: | 21,753: | -: -: | 31: 328 |
| Round Lake | 5: 5: | 43: 11: 307: | 19: 475: 78: | 5,960: 176: | 29,045: | 2: 13: | 22: 291 |
| | 40: 365: | 85: 1362: | 122: 4058: | 506: 23,391: | 980: 135,539: | 19: 125: | 210: 2,206 |
| Av. Volume | : 9: | : 16: | : 35: | : 46: | : 139: | : 7: | : 10 |
| | : .22: | : .6: | : 2.4: | : 14.0: | : 81.2: | : .1: | : 1.3 |
| Ratio | : .08: | : .5: | : .35: | : 2.0: | : 1.0: | : .25: | : 2.6 |

Total - 1962 trees 167,056 bd.ft.